

AD\_\_\_\_\_

AWARD NUMBER: W81XWH-04-1-0920

TITLE: Preclinical Evaluation of Serine/Threonine Kinase Inhibitors Against Prostate Cancer Metastases

PRINCIPAL INVESTIGATOR: Theresa Guise, M.D.

CONTRACTING ORGANIZATION: University of Virginia  
Charlottesville, Virginia 22904

REPORT DATE: November 2006

TYPE OF REPORT: Annual, Revised

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

| REPORT DOCUMENTATION PAGE   |             |                                   |                            | Form Approved<br>OMB No. 0704-0188                       |   |
|---|-------------|-----------------------------------|----------------------------|--|---|
| Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b> |             |                                   |                            |  |   |
| 1. REPORT DATE (DD-MM-YYYY)<br>01-11-2006   |             | 2. REPORT TYPE<br>Annual, Revised |                            | 3. DATES COVERED (From - To)<br>15 Oct 2005– 14 Oct 2006 |   |
| 4. TITLE AND SUBTITLE<br><br>Preclinical Evaluation of Serine/Threonine Kinase Inhibitors Against Prostate Cancer Metastases  |             |                                   |                            | 5a. CONTRACT NUMBER                                      |   |
|   |             |                                   |                            | 5b. GRANT NUMBER<br>W81XWH-04-1-0920                     |   |
|   |             |                                   |                            | 5c. PROGRAM ELEMENT NUMBER                               |   |
| 6. AUTHOR(S)<br><br>Theresa Guise, M.D.<br><br>E-Mail: <a href="mailto:tag4n@virginia.edu">tag4n@virginia.edu</a>   |             |                                   |                            | 5d. PROJECT NUMBER                                       |   |
|   |             |                                   |                            | 5e. TASK NUMBER  |   |
|   |             |                                   |                            | 5f. WORK UNIT NUMBER                                     |   |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)<br><br>University of Virginia<br>Charlottesville, Virginia 22904   |             |                                   |                            | 8. PERFORMING ORGANIZATION REPORT NUMBER                 |   |
| 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)<br>U.S. Army Medical Research and Materiel Command<br>Fort Detrick, Maryland 21702-5012   |             |                                   |                            | 10. SPONSOR/MONITOR'S ACRONYM(S)                         |   |
|   |             |                                   |                            | 11. SPONSOR/MONITOR'S REPORT NUMBER(S)                   |   |
| 12. DISTRIBUTION / AVAILABILITY STATEMENT<br>Approved for Public Release; Distribution Unlimited  |             |                                   |                            |  |   |
| 13. SUPPLEMENTARY NOTES   |             |                                   |                            |  |   |
| 14. ABSTRACT<br><br>A central tenet of the field of bone metastases is that the bone microenvironment supplies factors, such as TGF-beta, stimulating prostate cancer cell signaling and altering their phenotype. TGF-beta signaling in cancer is however complex and can lead to the activation of numerous genes. We have identified many of these genes by microarray analysis and have validated the gene reported here. Of these, PMEPA1 as the most highly upregulated gene. We have cloned the PMEPA1 promoter and full-length gene and have begun promoter analysis of the TGFβbetaresponse element. We are in the process of overexpressing PMEPA1 in prostate cancer lines. In vivo experiments are in progress to determine the effect of a TGFbeta RI kinase inhibitor, SD-208, on the development and progression of prostate cancer metastases to bone due to PC-3, LuCAP and C42B prostate cancers.                                       |             |                                   |                            |  |   |
| 15. SUBJECT TERMS<br>Bone metastases, prostate cancer, TGF-beta, PMEPA1   |             |                                   |                            |  |   |
| 16. SECURITY CLASSIFICATION OF:   |             |                                   | 17. LIMITATION OF ABSTRACT | 18. NUMBER OF PAGES                                      | 19a. NAME OF RESPONSIBLE PERSON           |
| a. REPORT   | b. ABSTRACT | c. THIS PAGE                      |                            |  | USAMRMC                                   |
| U   | U           | U                                 | UU                         | 21   | 19b. TELEPHONE NUMBER (include area code) |

## **Table of Contents**

|                     |            |
|---------------------|------------|
| Introduction        | Page 2     |
| Body                | Page 2-3   |
| Hypotheses          | Page 3     |
| Specific Aims       | Page 3-4   |
| Results             | Page 5-6   |
| Reportable Outcomes | Page 7-9   |
| Conclusions         | Page 10    |
| References          | Page 11-19 |

## Introduction

Prostate cancer has a propensity to grow in the skeleton and cause significant morbidity. Once housed in bone, prostate cancer is incurable. Bone is a rich storehouse of growth factors, which stimulate signaling in metastatic cancer cells. Bone-derived TGF $\beta$  increases tumor secretion of factors that activate bone remodeling, fueling a vicious cycle, which drives the growth and survival of prostate bone metastases. In prostate cancer cells, TGF $\beta$  signals through two receptor subunits and, further downstream, p38 MAP kinase. Hypothesis: *TGF $\beta$  mediates prostate cancer metastases to bone via p38 MAP kinase pathway. TGF $\beta$  and/or p38MAP kinase signaling inhibitors will reduce the development and progression of prostate cancer bone metastases to bone.* Two orally active inhibitors of these serine/threonine kinases will be tested in an animal model of prostate cancer bone metastases. We propose three Specific Aims. Aim 1: To test a TGF $\beta$ RI kinase inhibitor and a p38 MAPK inhibitor against three human prostate cancer models of skeletal metastasis in mice. Aim 2: To determine the molecular targets of these inhibitors in prostate cancer cells *in vitro* and test their impact on tumor growth and bone metastases *in vivo*. Aim 3: To test the efficacy of combined TGF $\beta$ RI and p38 MAP kinase inhibitors against three prostate cancer models *in vivo*.

## Body

**BACKGROUND.** The skeleton is a major site of metastasis by advanced prostate cancer. Last year 220,900 cases of prostate cancer were diagnosed in the United States, where it is now the most commonly diagnosed cancer and the second most common cause of cancer mortality in men, with 28,900 deaths (Crawford, 2003). One fourth of diagnosed patients will die from the disease, the majority of them with metastases to the skeleton. Once cancer becomes housed in bone, it is incurable. The average survival from time of diagnosis of skeletal metastases in prostate cancer patients is 40 months. When prostate tumor cells metastasize to the skeleton, the most common response is osteoblastic: characterized by net formation of disorganized new bone, which results in fractures, severe and intractable bone pain, and nerve compression. Metastasis to bone thus causes prolonged, serious morbidity for many prostate cancer patients. Treatment to prevent or halt the progression of bone metastases (Reddi et al, 2003; O’Keefe and Guise, 2003). would increase survival and improve quality of life for men with prostate cancer

***Transforming growth factor- $\beta$  in cancer is a two-edged sword.*** TGF $\beta$  is a growth inhibitor and a tumor suppressor at early stages of the oncogenic cascade. However, advanced cancers often lose the growth inhibition by TGF $\beta$  but continue to respond to the factor. The net effect is that TGF $\beta$  is a metastasis enhancer for advanced cancers. Since bone is a major source of active TGF $\beta$ , the factor plays a crucial role in the vicious cycle of bone metastases. Blockade of the TGF $\beta$  pathway effectively decreases metastases in several animal models (Yin et al, 1996; Muraoka et al, 2002; Yang et al, 2002).

***Transforming growth factor- $\beta$  in bone*** is released from mineralized matrix in active form by osteoclastic resorption (Dallas et al, 2002), which is very prominent in prostate cancer metastases. TGF $\beta$  acts on tumor cells to increase the secretion of factors that inappropriately stimulate bone cells (Chirgwin & Guise, 2003a,b). The interactions

between bone and cancer constitute a vicious cycle, which enhances skeletal metastases (Mundy, 2002). Extensive data show that TGF $\beta$  is a major bone-derived factor responsible for driving the vicious cycle of cancer metastases in bone. TGF $\beta$  increases tumor secretion of factors such as endothelin-1, IL-6, IL-11, PTHrP, and VEGF. These factors stimulate both osteoblastic synthesis of disorganized new bone and osteolytic destruction of the skeleton adjacent to tumor cells. The cellular and molecular components of the vicious cycle between tumor and bone offer opportunities for therapeutic intervention to decrease skeletal metastases (Coleman, 2002; Guise & Chirgwin, 2003a). TGF $\beta$  in particular is an important target for intervention against prostate cancer skeletal metastases.

***Therapy to block TGF $\beta$  signaling in bone metastases.*** Previous work has demonstrated the effectiveness of TGF-beta inhibition to decrease metastases, but these experiments have used protein-based treatment or ex vivo manipulations of the tumor cells (Yin et al, 1996; Muraoka et al, 2002; Yang et al, 2002). Orally active small-molecule inhibitors of the TGF $\beta$  pathway would be much more practical. This proposal will test two inhibitors of serine/threonine kinases. The first directly targets the TGF $\beta$  receptor kinase. The second targets p38 MAP kinase, which is a major downstream effector of TGF $\beta$  signaling in cancer cells. Both targets are serine/threonine kinases. Our preliminary data show that inhibition of TGF $\beta$  signaling is effective in an animal model of cancer bone metastases. The work proposed will test the two serine/threonine kinases inhibitors in animal models of human prostate cancer in bone: one in which the response is osteolytic, two others in which it is osteoblastic. The experiments proposed will rapidly provide the preclinical data necessary for these two drugs to be placed in clinical trials for prostate cancer bone metastases.

**Hypotheses:** 1) TGF $\beta$  mediates prostate cancer metastases to bone via p38 MAP kinase. Specific serine/threonine kinase small-molecule inhibitors of the type I TGF $\beta$  receptor kinase and of p38 MAP kinase will reduce the development and progression of prostate cancer metastases to bone, due to either osteoblastic or osteolytic diseases. 2) Orally active inhibitors of these serine/threonine kinases will be effective in animal models of prostate cancer bone metastases to decrease metastases and tumor burden and to increase survival. 3) The two drugs may be more effective in combination than singly, if p38 MAP kinase also mediates TGF $\beta$ -independent metastatic functions. 4) Specific targets of TGF $\beta$  signaling in prostate cancer cells contribute directly to the bone phenotype of metastases. One such factor may be the type I membrane protein PMEPA1, which is regulated by TGF $\beta$  and expressed by prostate cancers. 5) Expression of PMEPA1 on the surface of cancer cells will increase the development and progression of prostate cancer metastases to bone.

**Specific Aim 1:** To determine the effect of TGF $\beta$ RI kinase or p38 MAPK blockade separately against 3 human prostate cancer models of skeletal metastasis in mice (hypotheses 1 & 2).

**Results and Progress:** This experiment is in the final stages of progress with respect to the TGF $\beta$ RI kinase inhibitor and the p38MAP kinase inhibitor. We have tested the effects of the TGF $\beta$ RI kinase, SD-208, on the development and progression of bone

metastases due to PC-3, LuCAP23.1 and C42B prostate cancer. This aim has taken longer than originally planned because we had to determine long-term pharmacokinetics for drug delivery in the food. We have pharmacokinetic data that 50 and 100 mg/kg of SD-208 added to food result in drug levels that were effective in our mouse model of breast cancer metastases to bone. Here we show data that TGF $\beta$ RI kinase, SD-208, reduced osteolytic bone metastases due to PC-3, but increased osteoblastic bone metastases due to LuCAP23.1. There was no effect on the mixed tumor, C42B, which is unresponsive to TGF $\beta$ . We observed different effects of p38MAP kinase inhibition in these models, which are described below. The p38MAP kinase inhibitor, SD-282 increased bone metastases due to PC-3 prostate cancer and had no effect on LuCAP23.1 or C42B.

**Specific Aim 2:** To determine the molecular targets of the inhibitors in prostate cancer cells in vitro by gene array analysis (hypothesis 4). The role of an already-identified target of TGF $\beta$ , PMEPA1, will be tested in the animal models by overexpressing it in 2 prostate cancer cell lines (hypothesis 5).


*Results and Progress:* The majority of the work from year 1 has been performed on this aim. Gene array targets of TGF $\beta$  on PC-3 prostate cancer were validated by quantitative real-time PCR and were described in the progress report for year one. We are currently performing stable knockdown of PMEPA1 as well as overexpression and plan to test stable cell lines in the bone metastases model. We have also made deletion constructs of the promoter and are in the process of studying the TGF $\beta$  regulation of this promoter, which contains a number of Smad binding sites. Finally, we have made mutations of the PPXY sites of PMEPA1, which could bind to inhibitors of the TGF $\beta$  pathway. These mutants will be expressed in PC-3 prostate cancer and studied for effects on TGF $\beta$  signaling and bone metastases.

**Specific Aim 3:** To test the efficacy of combined T $\beta$ RI and p38 MAPK inhibitors against 3 prostate cancer models in vivo (hypothesis 3).

*Results and Progress:* This aim has not been started and is planned for the last year of the proposal.

## Results:


*p38MAP kinase inhibitor increases osteolytic bone metastases due to PC-3 prostate cancer.* Most recently, we found that treatment with a p38MAP kinase inhibitor, SD-282, accelerated development of osteolytic lesions due to PC-3 prostate cancer (**Figure 1**). In this experiment, mice were treated with SD-282 when bone metastases were identified on radiographs, at 4 weeks. Doses used were based on pharmacokinetic studies done by the manufacturer, Scios, Inc. We found similar results using the osteolytic breast cancer model, MDA-MB-231: p38MAP kinase inhibition increased osteolytic bone metastases. We have ongoing experiments testing this inhibitor on bone metastases due to LuCAP23.1 and C42B. To




osteolytic bone metastases and improved survival in mice bearing PC-3 prostate cancers (both treatment and prevention protocols). (**Figure 2, 3**) Taken together with data that a p38MAPK inhibitor was ineffective and even increased PC-3 and MDA-MB-231 bone metastases, it may be better to target the Smad pathway than total TGF $\beta$  signaling. The latter may be less effective, if downstream p38MAP kinase blockade adversely affects bone metastases.

date, there has been no effect of SD-282 on either group. In this, and in the PC-3 experiment, quantitative bone histomorphometry will be performed.

*TGF $\beta$ RI kinase inhibitor reduces osteolytic bone metastases due to PC-3 prostate cancer.* In contrast, and similar to results observed with osteolytic breast cancer model, MDA-MB-231, the TGF $\beta$ RI kinase inhibitor, SD-208, reduced







## 2. TGFβRI kinase inhibitor accelerates osteoblastic bone metastases due to prostate cancer LuCAP23.1.

Since our prior studies included only osteolytic bone metastases models, and TGFβ has been implicated in the pathogenesis of prostate cancer metastases to bone, we tested SD-208 in a model of human prostate cancer, LuCAP23.1, which grows as osteoblastic lesions when directly injected into bone. LuCAP23.1 (obtained from our collaborator, Robert Vessella, University of Washington) is an androgen sensitive, PSA-producing human tumor derived from an osteoblastic bone metastasis. It causes osteoblastic lesions in 10-12 weeks. **Figure 4** shows representative radiographs from LuCAP23.1 bearing mice, treated with SD-208 at 20 or 60 mg/kg/day after inoculation into bone. Drug was started at 12 weeks, after lesions were evident by radiographs. The experiment, still in progress, shows clearly that osteoblastic bone metastases are accelerated by treatment with SD-208. These preliminary data indicate that TGFβRI kinase inhibition may worsen osteoblastic metastases. Since our preliminary data also indicate that SD-208 has direct effects to increase osteoblast activity, this host response may accelerate osteoblastic disease in prostate cancer. Since LuCAP23.1 is a xenograft which

cannot be studied in culture as a cell line, we will test TGFβ inhibition on other osteoblastic models of breast and prostate cancer. ZR-75-1 is a human breast cancer which causes osteoblastic metastases due to tumor production of ET-1 (Yin et al 2003). We recently derived a variant of the PC-3 prostate cancer line that causes osteoblastic bone metastases, PC-3OB, and are now characterizing it (**Figure 5**). We are also testing SD-208 on bone metastases due to the mixed osteolytic & osteoblastic tumor C42B and, thus far, there has been no effect observed, compared to control. Quantitative histomorphometry will be performed on bones from all experiments.





## Reportable Outcomes

### Presentations:

- Targeting the endothelin axis in osteoblastic bone metastases: mechanisms and implications. Prostate Cancer Foundation Scientific Retreat, Scottsdale, AZ, Oct 2005.
- What makes bone a favorable site for metastases? AACR Tumor Microenvironment and Protease meeting, Bonita Springs, FL, Dec 2005.
- Bisphosphonate update 2006. 26<sup>th</sup> Annual Scripps Clinical Hematology and Oncology Meeting, San Diego, CA, Feb 2006.
- TGF $\beta$  in bone metastases: implications for therapy. Research Seminar, University of Lyon, Lyon France, Mar 2006.
- TGF $\beta$  in bone metastases: pathophysiology to treatment. Research Seminar, INSERM Unit 627, Hospital St. Louis, Paris, France, Mar 2006.
- Gene signatures in bone metastases: Role of TGF $\beta$ . European Calcified Tissue Society Meeting, Prague, Czech Republic, May 2006.
- TGF $\beta$  in skeletal complications of malignancy. Seminar at Schering, Berlin, Germany, May 2006.
- Osteoporosis in the cancer patient. First international meeting on secondary causes of osteoporosis, Florence, Italy, Jul 2006.
- PTHrP in osteoblastic bone metastases. PPP meeting. Newfoundland, Canada, Jul 2006.
- Pathophysiology of metastases. Regional Medical Liaison meeting, UCSF, San Francisco, CA, Jul 2006.
- Bone micrometastases. International Breast Cancer Conference. Kona, HI, Aug 2006.
- Pathophysiology of Bone Metastases. International Metastases Meeting, Tokushima, Japan, Sep 2006.
- TGF $\beta$  in bone metastases. Animal Model Working Group, American Society for Bone and Mineral Research Meeting, Philadelphia, PA, Sep 2006.
- Molecular mechanisms of bone metastases: Role of TGF $\beta$ . Fat and Bone Meeting, Madrid, Spain, Sep 2006.
- Molecular mechanisms of bone metastases: osteolytic and osteoblastic. Italian Cancer Society Meeting, Bari, Italy, Oct 2006.
- TGF $\beta$  in bone metastases: pathophysiology to treatment. Visiting Professor, University of Minnesota, Dec 2006.
- TGF $\beta$  signaling in breast cancer bone metastases: friend or foe? Cancer and Bone Society Meeting, San Antonio, TX Dec 2006.
- Molecular mechanisms of bone metastases: insight into therapy. Endocrine Grand Rounds, University of Texas Health Science Center at San Antonio, TX, Dec 2006.
- Skeletal complications of cancer and cancer treatment. Bone Club, San Antonio, TX, Dec 2006.
- Skeletal health in the cancer patient. Maine State Osteoporosis Meeting, Sugarloaf, MN, Jan 2007.
- Effects of bisphosphonates on tumor cells. Consensus on Bone Loss in Cancer Patients on Aromatase Inhibitors. Geneva, Switzerland, February, 2007.
- TGF $\beta$  in bone metastases: pathophysiology to treatment. Institute for Molecular Medicine, University of Lisbon, Lisbon, Portugal, March 2007
- RANK Ligand in prostate cancer metastases to bone. Medical Grand Rounds, Hospital Santa Ana, University of Lisbon, Lisbon, Portugal, March 2007
- Skeletal health in the cancer patient. Endocrine Grand Rounds, Oregon Health Sciences University, Portland, OR. April 2007

- TGF $\beta$  signaling in breast cancer bone metastases: friend or foe? Research Seminar, Oregon Health Sciences University, Portland, OR. April 2007
- TGF $\beta$  in bone metastases: pathophysiology to treatment. Endocrinology Grand Rounds, Mount Sinai School of Medicine, New York, NY, April 2007
- TGF $\beta$  signaling in breast cancer bone metastases: friend or foe? Advances in Mineral Metabolism Meeting, Snowmass, CO, April, 2007
- TGF $\beta$  signaling in cancer metastases to bone: friend or foe? Cleveland Clinic, April, 2007
- Biology of bone metastases. FASEB Meeting, Washington, D.C., April 2007
- Endothelins in pathologic and normal bone remodeling. Bone Club. University of Pittsburgh, Pittsburgh, PA, May 2007
- Endothelins in pathologic and normal bone remodeling. Research Seminar, Wyeth, Collegeville, PA. May 2007

#### Publications

- **Guisse TA.** Do markers of bone turnover predict clinical outcome in patients with bone metastases? Nat Clin Pract Endocrinol Metab. 2(5):258-9, May 2006. PMID: 16932296.
- Seth P, Wang ZG, Pister A, Zafar MB, Kim S, **Guisse T**, Wakefield L. Development of oncolytic adenovirus armed with a fusion of soluble transforming growth factor-beta receptor II and human immunoglobulin Fc for breast cancer therapy. Hum Gene Ther. 17(11):1152-60, Nov 2006. PMID: 17032151.
- Theriault RL, Biermann JS, Brown E, Brufsky A, Demers L, Grewal RK, **Guisse T**, Jackson R, McEnery K, Podoloff D, Ravdin P, Shapiro CL, Smith M, Van Poznak CH. NCCN Task Force Report: Bone health and cancer care. J Natl Compr Canc Netw. 4 Suppl 2:S1-20, May 2006. PMID: 16737674.
- Clines GA, Mohammad KS, Bao Y, Stephens OW, Suva LJ, Shaughnessy JD Jr, Fox JW, Chirgwin JM, **Guisse TA.** Dickkopf homolog 1 mediates endothelin-1-stimulated new bone formation. Mol Endocrinol. 21(2):486-98, Feb 2007. Epub 2006 Oct 26. PMID: 17068196.
- **Guisse TA**, Mohammad KS, Clines G, Stebbins EG, Wong DH, Higgins LS, Vessella R, Corey E, Padalecki S, Suva L, Chirgwin JM. Basic mechanisms responsible for osteolytic and osteoblastic bone metastases. Clin Cancer Res. 12(20):6213s-6s, Oct 2006. PMID: 17062703.
- Lipton A, Berenson JR, Body JJ, Boyce BF, Bruland OS, Carducci MA, Cleeland CS, Clohisey DR, Coleman RE, Cook RJ, **Guisse TA**, Pearse RN, Powles TJ, Rogers MJ, Roodman GD, Smith MR, Suva LJ, Vessella RL, Weilbaecher KN, King L. Advances in treating metastatic bone cancer: summary statement for the First Cambridge Conference. Clin Cancer Res. 12(20):6209s-12s, Oct 2006. PMID: 17062702.
- **Guisse TA.** Bone loss and fracture risk associated with cancer therapy. Oncologist. 11(10):1121-1131, Nov/Dec 2006. PMID: 17110632.
- Javelaud D, Mohammad KS, McKenna CR, Fournier P, Luciani F, Niewolna M, André J, Delmas V, Larue L, **Guisse TA**, Mauviel A. Stable overexpression of Smad7 in human melanoma cells impairs bone metastasis. Cancer Res. 67(5):2317-24, Mar 2007. PMID: 17332363.

- Brown SA, **Guise TA**. Drug insight: the use of bisphosphonates for the prevention and treatment of osteoporosis in men. *Nat Clin Pract Urol*. 2007 Jun;4(6):310-20. PMID: 17551535
- Kingsley LA, Chirgwin JM, **Guise TA**. Breaking new ground to build bone. *Proc Natl Acad Sci USA* In Press June 2007
- Chirgwin JM, **Guise TA**. Does prostate-specific antigen contribute to bone metastases? *Clin Cancer Res*. 12(5):1395-7, Mar 2006. PMID: 16533760.
- Heras-Herzig A, **Guise TA**. Disorders of calcium metabolism. IN: *Therapy of Renal Diseases and Related Disorders*, 4th edition. WN Suki, SG Massry (eds). Kluwer Academic Publishers, Chapter 5, In Press, 2006.
- Clines GA, **Guise TA**. Mechanisms of bone destruction and bone formation by metastatic tumors. *Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism*, sixth edition, edited by Murray J. Favus, MD et al. In Press 2006.
- Fournier P, Chirgwin JM, **Guise TA**. New insights into the role of T cells in the vicious cycle of bone metastases. *Curr Opin Rheumatol*. 18(4):396-404, Jul 2006. PMID: 16763461.
- Siclari VA, **Guise TA**, Chirgwin JM. Molecular interactions between breast cancer cells and the bone microenvironment drive skeletal metastases. *Cancer Metastasis Rev*. 25(4):621-33, Dec 2006. PMID: 17165131.
- Heras-Herzig A, Kozlow W, Brown S, **Guise TA**. Osteoporosis associated with cancer and cancer treatment. Chapter IN: *Osteoporosis* 3rd Ed. Academic Press. Editors: Marcus, Feldman, Rosen, Nelson; In Press 2007.

## Conclusions

A central tenet of the field of bone metastases is that the bone microenvironment supplies factors, such as TGF- $\beta$ , stimulating prostate cancer cell signaling and altering their phenotype. TGF- $\beta$  signaling in cancer is however complex and can lead to the activation of numerous genes. We have identified many of these genes by microarray analysis and have validated the gene reported here. Of these, PMEPA1 as the most highly upregulated gene. We have cloned the PMEPA1 promoter and full-length gene and have begun promoter analysis of the TGF $\beta$  response element. We are in the process of overexpressing and silencing PMEPA1 in prostate cancer lines.

In vivo experiments to determine the effect of a TGF $\beta$  RI kinase inhibitor, SD-208, on the development and progression of prostate cancer metastases to bone due to PC-3, LuCAP and C42B prostate cancers have shown differential effects, depending on the radiographic phenotype of the bone metastases. SD-208 improves osteolytic bone metastases due to PC-3, but increases osteoblastic bone metastases due to LuCAP23.1 and has no effect on mixed C42B lesion. The results in C42B are not surprising, given the fact that this line is unresponsive to TGF $\beta$ . However, the effect of this compound to increase osteoblastic bone metastases is a significant concern. This needs to be confirmed the information utilized with respect to design of future clinical trials. We have initiated an agreement with Eli Lilly to study another TGF $\beta$ RI kinase inhibitor which is currently in clinical trials for patients with bone metastases due to all solid tumors.

## References

Abasolo I, Yang L, Haleem R, Xiao W, Pio R, Cuttitta F, Montuenga LM, Kozlowski JM, Calvo A, Wang Z. Overexpression of adrenomedullin gene markedly inhibits proliferation of PC3 prostate cancer cells in vitro and in vivo. *Mol Cell Endocrinol*, 199:179-187, 2003.

Adler HL, McCurdy MA, Kattan MW, Timme TL, Scardino PT, Thompson TC. Elevated levels of circulating interleukin-6 and transforming growth factor-beta1 in patients with metastatic prostatic carcinoma. *J Urol*, 161:182-187, 1999.

Akhurst RJ, Derynck R. TGF-beta signaling in cancer--a double-edged sword. *Trends Cell Biol*, 11:S44-51, 2001.

Anan T, Nagata Y, Koga H, Honda Y, Yabuki N, Miyamoto C, Kuwano A, Matsuda I, Endo F, Saya H, Nakao M. Human ubiquitin-protein ligase Nedd4: expression, subcellular localization and selective interaction with ubiquitin-conjugating enzymes. *Genes Cells*, 3:751-763, 1998.

Bakin RE, Gioeli D, Sikes RA, Bissonette EA, Weber MJ. Constitutive activation of the Ras/mitogen-activated protein kinase signaling pathway promotes androgen hypersensitivity in LNCaP prostate cancer cells. *Cancer Res*, 63:1981-1989, 2003a.

Bakin RE, Gioeli D, Bissonette EA, Weber MJ. Attenuation of Ras signaling restores androgen sensitivity to hormone-refractory C4-2 prostate cancer cells. *Cancer Res*, 63:1975-1980, 2003b.

Bartholin L, Wessner LL, Chirgwin JM, Guise TA. The human Cyr61 gene is a transcriptional target of transforming growth factor beta in cancer cells. *Cancer Lett*. 2006

Bello-DeOcampo D, Tindall DJ. TGF-beta1/Smad signaling in prostate cancer. *Curr Drug Targets*, 4:197-207. 2003.

Bendre MS, Gaddy-Kurten D, Mon-Foote T, Akel NS, Skinner RA, Nicholas RW, Suva LJ. Expression of interleukin 8 and not parathyroid hormone-related protein by human breast cancer cells correlates with bone metastasis in vivo. *Cancer Res*, 62:5571-5579, 2002.

Blackledge G. Growth Factor Receptor Tyrosine Kinase Inhibitors; Clinical Development and Potential for Prostate Cancer Therapy. *J Urol*, 170:S77-S83, 2003.

Blanchere M, Saunier E, Mestayer C, Broshuis M, Mowszowicz I. Alterations of expression and regulation of transforming growth factor beta in human cancer prostate cell lines. *J Steroid Biochem Mol Biol*, 82:297-304, 2002.

Boyde A, Maconnachie E, Reid SA, Delling G, Mundy GR. Scanning electron microscopy in bone pathology: Review of methods. Potential and applications. *Scanning Electron Microscopy IV*:1537-1554, 1986.

Brigstock DR. Regulation of angiogenesis and endothelial cell function by connective tissue growth factor (CTGF) and cysteine-rich 61 (CYR61). *Angiogenesis*, 5:153-165, 2002.

Brunschwig EB, Wilson K, Mack D, Dawson D, Lawrence E, Willson JK, Lu S, Nosrati A, Rerko RM, Swinler S, Beard L, Lutterbaugh JD, Willis J, Platzer P, Markowitz S. PMEPA1, a transforming growth factor-beta-induced marker of terminal colonocyte differentiation whose expression is maintained in primary and metastatic colon cancer. *Cancer Res*, 63:1568-1575, 2003.

Carducci MA, Padley RJ, Breul J, Vogelzang NJ, Zonnenberg BA, Daliani DD, Schulman CC, Nabulsi AA, Humerickhouse RA, Weinberg MA, Schmitt JL, Nelson JB. Effect of endothelin-A receptor blockade with atrasentan on tumor progression in men with hormone-refractory prostate cancer: a randomized, phase II, placebo-controlled trial. *J Clin Oncol*, 21:679-689, 2003.

Cher ML. Mechanisms governing bone metastasis in prostate cancer. *Curr Opin Urol*, 11:483-488, 2001.

Chesneau V, Becherer JD, Zheng Y, Erdjument-Bromage H, Tempst P, Blobel CP. Catalytic properties of ADAM19. *J Biol Chem*, 278:22331-22340, 2003.

Chipuk JE, Cornelius SC, Pultz NJ, Jorgensen JS, Bonham MJ, Kim SJ, Danielpour D. The androgen receptor represses transforming growth factor-beta signaling through interaction with Smad3. *J Biol Chem*, 277:1240-1248, 2002.

Chirgwin JM, Guise TA. Molecular mechanisms of tumor-bone interactions in osteolytic metastases. *Crit Rev Eukaryot Gene Expr*, 10:159-178, 2000.

Chirgwin JM, Guise TA. Molecular mechanisms of cancer metastases to bone. *Curr Opin Orthop*, 14:317-321, 2003a.

Chirgwin JM, Guise TA. Cancer metastasis to bone. *Science & Medicine*, 9:140-151, 2003b.

Chung LW. Prostate carcinoma bone-stroma interaction and its biologic and therapeutic implications. *Cancer*, 97(3 Suppl):772-778, 2003.

Corey E, Quinn JE, Bladou F, Brown LG, Roudier MP, Brown JM, Buhler KR, Vessella RL. Establishment and characterization of osseous prostate cancer models: intra-tibial injection of human prostate cancer cells. *Prostate*, 52:20-33, 2002.

- Crawford ED. Epidemiology of prostate cancer. *Urology*, 62(6 Suppl 1):3-12, 2003.
- Crawford *et al.* 1998. *Cell*. 93:1159
- Good *et al.* 1990. *PNAS*. 87:6624
- Volpert *et al.* 1998. *PNAS*. 95:6343
- Filleur *et al.* 2001. *Genes & Dev*. 15:1373
- Fontana *et al.* 2005. *Int. J. Cancer*. 116:686
- Dallas SL, Rosser JL, Mundy GR, Bonewald LF. Proteolysis of latent transforming growth factor-beta (TGF-beta)-binding protein-1 by osteoclasts. A cellular mechanism for release of TGF-beta from bone matrix. *J Biol Chem*, 277:21352-21360, 2002.
- Derynck R, Zhang YE. Smad-dependent and Smad-independent pathways in TGF-beta family signalling. *Nature*, 425:577-584, 2003.
- Deftos LJ. Prostate carcinoma: production of bioactive factors. *Cancer*, 88(12 Suppl):3002-3008, 2000.
- Feng XH, Liang YY, Liang M, Zhai W, Lin X. Direct interaction of c-Myc with Smad2 and Smad3 to inhibit TGF-beta-mediated induction of the CDK inhibitor p15(Ink4B). *Mol Cell*, 9:133-143, 2002.
- Festuccia C, Angelucci A, Gravina GL, Villanova I, Teti A, Albini A, Bologna M, Abini A. Osteoblast-derived TGF-beta1 modulates matrix degrading protease expression and activity in prostate cancer cells. *Int J Cancer*, 85:407-415, 2000.
- Fidler IJ. The pathogenesis of cancer metastasis: the 'seed and soil' hypothesis revisited. *Nat Rev Cancer*, 3:1-6, 2003.
- Fu Z, Smith PC, Zhang L, Rubin MA, Dunn RL, Yao Z, Keller ET. Effects of raf kinase inhibitor protein expression on suppression of prostate cancer metastasis. *J Natl Cancer Inst*, 95:878-889, 2003a.
- Fu Y, O'Connor LM, Shepherd TG, Nachtigal MW. The p38 MAPK inhibitor, PD169316, inhibits transforming growth factor beta-induced Smad signaling in human ovarian cancer cells. *Biochem Biophys Res Commun*, 310:391-397, 2003b.
- Gallwitz WE, Guise TA, Mundy GR. Guanosine nucleotides inhibit different syndromes of PTHrP excess caused by human cancers in vivo. *J Clin Invest*, 110:1559-1572, 2002.
- Giannini G, Ambrosini MI, Di Marcotullio L, Cerignoli F, Zani M, MacKay AR, Screpanti I, Frati L, Gulino A. EGF- and cell-cycle-regulated STAG1/PMEPA1/ERG1.2

belongs to a conserved gene family and is overexpressed and amplified in breast and ovarian cancer. *Mol Carcinog*, 8:188-200, 2003.

Gioeli D, Mandell JW, Petroni GR, Frierson HF Jr, Weber MJ. Activation of mitogen-activated protein kinase associated with prostate cancer progression. *Cancer Res*, 59:279-284, 1999.

Gioeli D, Zecevic M, Weber MJ. Immunostaining for activated extracellular signal-regulated kinases in cells and tissues. *Methods Enzymol*, 332:343-353, 2001.

Granchi S, Brocchi S, Bonaccorsi L, Baldi E, Vinci MC, Forti G, Serio M, Maggi M. Endothelin-1 production by prostate cancer cell lines is up-regulated by factors involved in cancer progression and down-regulated by androgens. *Prostate*, 49:267-277, 2001.

Grotendorst GR, Okochi H, Hayashi N. A novel transforming growth factor beta response element controls the expression of the connective tissue growth factor gene. *Cell Growth Differ*, 7:469-480, 1996.

Guise TA, Yin JJ, Taylor SD, Kumagai Y, Dallas M, Boyce BF, Yoneda T, Mundy GR. Evidence for a causal role of parathyroid hormone-related protein in the pathogenesis of human breast cancer-mediated osteolysis. *J Clin Invest*, 98:1544-1549, 1996.

Guise TA, Mundy GR. Cancer and bone. *Endocr Rev*, 19:18-55, 1998.

Guise TA, Chirgwin JM. Role of bisphosphonates in prostate cancer bone metastases. *Semin Oncol*, 30:717-723, 2003a.

Guise TA, Chirgwin JM. Transforming growth factor-beta in osteolytic breast cancer bone metastases. *Clin Orthop*, 415:S32-38, 2003b.

Hauschka PV, Mavrakos AE, Iafrati MD, Doleman SE, Klagsbrun M. Growth factors in bone matrix. Isolation of multiple types by affinity chromatography on heparin-Sepharose. *J Biol Chem*, 261:12665-12674, 1986.

Hayes SA, Huang X, Kambhampati S, Platanias LC, Bergan RC. p38 MAP kinase modulates Smad-dependent changes in human prostate cell adhesion. *Oncogene*, 22:4841-4850, 2003.

Hortobagyi GN, Theriault RL, Porter L, Blayney D, Lipton A, Sinoff C, Wheeler H, Simeone JF, Seaman J, Knight RD. Efficacy of pamidronate in reducing skeletal complications in patients with breast cancer and lytic bone metastases. Protocol 19 Aredia Breast Cancer Study Group. *N Engl J Med*, 335: 1785-1791, 1996.

Hwa V, Oh Y, Rosenfeld RG. Insulin-like growth factor binding protein-3 and -5 are regulated by transforming growth factor-beta and retinoic acid in the human prostate adenocarcinoma cell line PC-3. *Endocrine*, 6:235-242, 1997.



Itoh S, Thorikay M, Kowanetz M, Moustakas A, Itoh F, Heldin CH, ten Dijke P. Elucidation of Smad requirement in transforming growth factor-beta type I receptor-induced responses. *J Biol Chem*, 278:3751-3761, 2003.

Janda E, Lehmann K, Killisch I, Jechlinger M, Herzig M, Downward J, Beug H, Grunert S. Ras and TGF[beta] cooperatively regulate epithelial cell plasticity and metastasis: dissection of Ras signaling pathways. *J Cell Biol*, 156:299-313, 2002.

Johnson GL, Lapadat R. Mitogen-activated protein kinase pathways mediated by ERK, JNK, and p38 protein kinases. *Science*, 298:1911-1912, 2002.

Kakonen SM, Selander KS, Chirgwin JM, Yin JJ, Burns S, Rankin WA, Grubbs BG, Dallas M, Cui Y, Guise TA. Transforming growth factor-beta stimulates parathyroid hormone-related protein and osteolytic metastases via Smad and mitogen-activated protein kinase signaling pathways. *J Biol Chem*, 277:24571-24578, 2002.

Kang Y, Siegel PM, Shu W, Drobnjak M, Kakonen SM, Cordon-Cardo C, Guise TA, Massague J. A multigenic program mediating breast cancer metastasis to bone. *Cancer Cell*, 3:537-549, 2003.

Keller ET. The role of osteoclastic activity in prostate cancer skeletal metastases. *Drugs Today*, 38:91-102, 2002.

Kozawa O, Kawamura H, Hatakeyama D, Matsuno H, Uematsu T. Endothelin-1 induces vascular endothelial growth factor synthesis in osteoblasts: involvement of p38 mitogen-activated protein kinase. *Cell Signal*, 2:375-380, 2000.

Liao Y, Hung MC. Regulation of the activity of p38 mitogen-activated protein kinase by Akt in cancer and adenoviral protein E1A-mediated sensitization to apoptosis. *Mol Cell Biol*, 23:6836-6848, 2003.

Lum *et al.* 1999. *JBC*. 274:13613

Lynch CC, Hikosaka A, Acuff HB, Martin MD, Kawai N, Singh RK, Vargo-Gogola TC, Begtrup JL, Peterson TE, Fingleton B, Shirai T, Matrisian LM, Futakuchi M. MMP-7 promotes prostate cancer-induced osteolysis via the solubilization of RANKL. *Cancer Cell*. 2005 May;7(5):485-96.

Mackie EJ, Ramsey S. Modulation of osteoblast behaviour by tenascin. *J Cell Sci*, 109:1597-1604, 1996.

Mackie EJ, Abraham LA, Taylor SL, Tucker RP, Murphy LI. Regulation of tenascin-C expression in bone cells by transforming growth factor-beta. *Bone*, 22:301-307, 1998.

Maeda S, Hayashi M, Komiya S, Imamura T, Miyazono K. Endogenous TGF-beta signaling suppresses maturation of osteoblastic mesenchymal cells. *EMBO J*. 2004 Jan 29 [Epub ahead of print]

Merrell M, Suarez-Cuervo C, Harris KW, Vaananen HK, Selander KS. Bisphosphonate induced growth inhibition of breast cancer cells is augmented by p38 inhibition. *Breast Cancer Res Treat*, 81:231-241, 2003.

Miyake H, Pollak M, Gleave ME. Castration-induced up-regulation of insulin-like growth factor binding protein-5 potentiates insulin-like growth factor-I activity and accelerates progression to androgen independence in prostate cancer models. *Cancer Res*, 60:3058-3064, 2000.

Mohammad KS, Guise TA. Mechanisms of osteoblastic metastases: role of endothelin-1. *Clin Orthop*, 415:S67-74, 2003.

Mundy GR. Metastasis to bone: causes, consequences and therapeutic opportunities. *Nat Rev Cancer*, 2:584-593, 2002.

Muraoka RS, Dumont N, Ritter CA, Dugger TC, Brantley DM, Chen J, Easterly E, Roebuck LR, Ryan S, Gotwals PJ, Koteliansky V, Arteaga CL. Blockade of TGF-beta inhibits mammary tumor cell viability, migration, and metastases. *J Clin Invest*, 109:1551-1559, 2002.

Nelson JB. Endothelin Inhibition: Novel Therapy for Prostate Cancer. *J Urol*, 170:S65-S68, 2003.

O'Keefe RJ, Guise TA. Molecular mechanisms of bone metastasis and therapeutic implications. *Clin Orthop*, 415(Suppl):S100-104, 2003.

Park BJ, Park JI, Byun DS, Park JH, Chi SG. Oncogenic conversion of transforming growth factor-beta1 effect by oncogenic Ha-Ras-induced activation of the mitogen-activated protein kinase signaling pathway in human prostate cancer. *Cancer Res*, 60:3031-3038, 2000.

Park JI, Lee MG, Cho K, Park BJ, Chae KS, Byun DS, Ryu BK, Park YK, Chi SG. Transforming growth factor-beta1 activates interleukin-6 expression in prostate cancer cells through the synergistic collaboration of the Smad2, p38-NF-kappaB, JNK, and Ras signaling pathways. *Oncogene*, 22:4314-4332, 2003.

Pfitzenmaier J, Quinn JE, Odman AM, Zhang J, Keller ET, Vessella RL, Corey E. Characterization of C4-2 prostate cancer bone metastases and their response to castration. *J Bone Miner Res*, 18:1882-1888, 2003.

Pirtskhalaishvili G, Nelson JB. Endothelium-derived factors as paracrine mediators of prostate cancer progression. *Prostate*, 44:77-87, 2000.

Rae FK, Hooper JD, Nicol DL, Clements JA. Characterization of a novel gene, STAG1/PMEPA1, upregulated in renal cell carcinoma and other solid tumors. *Mol Carcinog*, 32:44-53, 2001.

Reddi AH, Roodman D, Freeman C, Mohla S: Mechanisms of tumor metastasis to the bone: challenges and opportunities. *J Bone Miner Res*, 18:190-194, 2003

Roberts AB, Wakefield LM. The two faces of transforming growth factor beta in carcinogenesis. *Proc Natl Acad Sci USA*, 100:8621-8623, 2003.

Ruggeri B, Singh J, Gingrich D, Angeles T, Albom M, Chang H, Robinson C, Hunter K, Dobrzanski P, Jones-Bolin S, Aimone L, Klein-Szanto A, Herbert JM, Bono F, Schaeffer P, Casellas P, Bourie B, Pili R, Isaacs J, Ator M, Hudkins R, Vaught J, Mallamo J, Dionne C. CEP-7055: a novel, orally active pan inhibitor of vascular endothelial growth factor receptor tyrosine kinases with potent antiangiogenic activity and antitumor efficacy in preclinical models. *Cancer Res*, 63:5978-5991, 2003.

Saad F, Gleason DM, Murray R, Tchekmedyian S, Venner P, Lacombe L, Chin JL, Vinholes JJ, Goas JA, Chen B. A randomized, placebo-controlled trial of zoledronic acid in patients with hormone-refractory metastatic prostate carcinoma. *J Natl Cancer Inst*, 94:1458-68, 2002.

Safadi FF, Xu J, Smock SL, Kanaan RA, Selim AH, Odgren PR, Marks SC Jr, Owen TA, Popoff SN.

Expression of connective tissue growth factor in bone: its role in osteoblast proliferation and differentiation in vitro and bone formation in vivo. *J Cell Physiol*, 196:51-62, 2003.

Shah AH, Tabayoyong WB, Kundu SD, Kim SJ, Van Parijs L, Liu VC, Kwon E, Greenberg NM, Lee C. Suppression of tumor metastasis by blockade of transforming growth factor beta signaling in bone marrow cells through a retroviral-mediated gene therapy in mice. *Cancer Res*, 62:7135-7138, 2002.

Schultz RM. Potential of p38 MAP kinase inhibitors in the treatment of cancer. *Prog Drug Res*, 60:59-92, 2003.

Shariat SF, Shalev M, Menesses-Diaz A, Kim IY, Kattan MW, Wheeler TM, Slawin KM. Preoperative plasma levels of transforming growth factor beta(1) (TGF-beta(1)) strongly predict progression in patients undergoing radical prostatectomy. *J Clin Oncol*, 19:2856-2864, 2001.

Siegel PM, Shu W, Cardiff RD, Muller WJ, Massague J. Transforming growth factor beta signaling impairs Neu-induced mammary tumorigenesis while promoting pulmonary metastasis. *Proc Natl Acad Sci USA*, 100:8430-8435, 2003.

Street J, Bao M, deGuzman L, Bunting S, Peale FV Jr, Ferrara N, Steinmetz H, Hoeffel J, Cleland JL, Daugherty A, van Bruggen N, Redmond HP, Carano RA, Filvaroff EH. Vascular endothelial growth factor stimulates bone repair by promoting angiogenesis and bone turnover. *Proc Natl Acad Sci U S A*. 2002 Jul 23;99(15):9656-61

Tang B, Vu M, Booker T, Santner SJ, Miller FR, Anver MR, Wakefield LM. TGF-beta switches from tumor suppressor to prometastatic factor in a model of breast cancer progression. *J Clin Invest*, 112:1116-1124, 2003.

Thalmann GN, Anezinis PE, Chang SM, Zhau HE, Kim EE, Hopwood VL, Pathak S, von Eschenbach AC, Chung LW. Androgen-independent cancer progression and bone metastasis in the LNCaP model of human prostate cancer. *Cancer Res*, 54:2577-2581, 1995.

Thomas RJ, Guise TA, Yin JJ, Elliott J, Horwood NJ, Martin TJ, Gillespie MT. Breast cancer cells interact with osteoblasts to support osteoclast formation. *Endocrinol*, 140:4451-4458, 1999.

Tokuda H, Hatakeyama D, Akamatsu S, Tanabe K, Yoshida M, Shibata T, Kozawa O. Involvement of MAP kinases in TGF-beta-stimulated vascular endothelial growth factor synthesis in osteoblasts. *Arch Biochem Biophys*, 415:117-125, 2003.

Tuxhorn JA, McAlhany SJ, Yang F, Dang TD, Rowley DR. Inhibition of transforming growth factor-beta activity decreases angiogenesis in a human prostate cancer-reactive stroma xenograft model. *Cancer Res*, 62:6021-6025, 2002.

Uy HL, Mundy GR, Boyce BF, Story BM, Dunstan CR, Yin JJ, Roodman GD, Guise TA. Tumor necrosis factor enhances parathyroid hormone-related protein-induced hypercalcemia and bone resorption without inhibiting bone formation in vivo. *Cancer Res*, 57:3194-3199, 1997.

van der Pluijm G, Sijmons B, Vloedgraven H, Deckers M, Papapoulos S, Lowik C. Monitoring metastatic behavior of human tumor cells in mice with species-specific polymerase chain reaction: elevated expression of angiogenesis and bone resorption stimulators by breast cancer in bone metastases. *J Bone Miner Res*, 16:1077-1091, 2001.

Wikstrom P, Damber J, Bergh A. Role of transforming growth factor-beta1 in prostate cancer. *Microsc Res Tech*, 52:411-419, 2001.

Wu TT, Sikes RA, Cui Q, Thalmann GN, Kao C, Murphy CF, Yang H, Zhau HE, Balian G, Chung LW. Establishing human prostate cancer cell xenografts in bone: induction of osteoblastic reaction by prostate-specific antigen-producing tumors in athymic and SCID/bg mice using LNCaP and lineage-derived metastatic sublines. *Int J Cancer*, 77:887-894, 1998.

Xu LL, Shanmugam N, Segawa T, Sesterhenn IA, McLeod DG, Moul JW, Srivastava S. A novel androgen-regulated gene, PMEPA1, located on chromosome 20q13 exhibits high level expression in prostate. *Genomics*, 66:257-263, 2000.

Xu LL, Shi Y, Petrovics G, Sun C, Makarem M, Zhang W, Sesterhenn IA, McLeod DG, Sun L, Moul JW, Srivastava S. PMEPA1, an androgen-regulated NEDD4-binding protein, exhibits cell growth inhibitory function and decreased expression during prostate cancer progression. *Cancer Res*, 63:4299-4304, 2003.

Yang YA, Dukhanina O, Tang B, Mamura M, Letterio JJ, MacGregor J, Patel SC, Khozin S, Liu ZY, Green J, Anver MR, Merlino G, Wakefield LM. Lifetime exposure to a soluble TGF-beta antagonist protects mice against metastasis without adverse side effects. *J Clin Invest*, 109:1607-1615, 2002.

Yin JJ, Selander K, Chirgwin JM, Dallas M, Grubbs BG, Wieser R, Massague J, Mundy GR, Guise TA. TGF-beta signaling blockade inhibits PTHrP secretion by breast cancer cells and bone metastases development. *J Clin Invest*, 103:197-206, 1999.

Yin JJ, Mohammad KS, Kakonen SM, Harris S, Wu-Wong JR, Wessale JL, Padley RJ, Garrett IR, Chirgwin JM, Guise TA. A causal role for endothelin-1 in the pathogenesis of osteoblastic bone metastases. *Proc Natl Acad Sci USA*, 100:10954-10959, 2003

Yoneda T, Sasaki A, Dunstan C, Williams PJ, Bauss F, De Clerck YA, Mundy GR. Inhibition of osteolytic bone metastasis of breast cancer by combined treatment with the bisphosphonate ibandronate and tissue inhibitor of the matrix metalloproteinase-2. *J Clin Invest*, 99:2509-2517, 1997.

Zheng *et al.* 2004. *JBC*. 279:42898

Zhou W, Park I, Pins M, Kozlowski JM, Jovanovic B, Zhang J, Lee C, Illo K. Dual regulation of proliferation and growth arrest in prostatic stromal cells by transforming growth factor-beta1. *Endocrinology*, 144:4280-4284, 2003.